

SIM61 References for Citations

- Selbmann L, Hoog de GS, Zucconi L, Isola D, Ruisi S, Gerrits van den Ende AHG, Ruibal C, De Leo F, Urzi C and Onofri S (2008). Drought meets acid: three new genera in a dothidealean clade of extremotolerant fungi. *Studies in Mycology* **61**: 1–20.
- Zalar P, Gostinčar C, Hoog GS de, Uršič V, Sudhadham M and Gunde-Cimerman N (2008). Redefinition of *Aureobasidium pullulans* and its varieties. *Studies in Mycology* **61**: 21–38.
- Yurlova NA, Hoog GS de and Fedorova LG (2008). The influence of ortho- and para-diphenoloxidase substrates on pigment formation in black yeast-like fungi *Studies in Mycology* **61**: 39–49.
- Gostinčar C, Turk M, Trbuha T, Vaupotič T, Plemenitaš A and Gunde-Cimerman N (2008). Expression of fatty-acid-modifying enzymes in the halotolerant black yeast *Aureobasidium pullulans* (de Bary) G. Arnaud under salt stress. *Studies in Mycology* **61**: 51–59.
- Vaupotič T, Veranic P, Petrovič U, Gunde-Cimerman N and Plemenitaš A (2008). HMG-CoA reductase is regulated by environmental salinity and its activity is essential for halotolerance in halophilic fungi. *Studies in Mycology* **61**: 61–66.
- Plemenitaš A, Vaupotič T, Lenassi M, Kogej T and Gunde-Cimerman N (2008). Adaptation of extremely halotolerant black yeast *Hortaea werneckii* to increased osmolarity: a molecular perspective at a glance. *Studies in Mycology* **61**: 67–75.
- Bonifaz A, Badali H, Hoog GS de, Cruz M, Araiza J, Cruz MA, Fierro L and Ponce RM (2008). Tinea nigra by *Hortaea werneckii*, a report of 22 cases from Mexico. *Studies in Mycology* **61**: 77–82.
- Harutyunyan S, Muggia L and Grube M (2008). Black fungi in lichens from seasonally arid habitats. *Studies in Mycology* **61**: 83–90.
- Gorbushina AA, Kotlova ER and Sherstneva OA (2008). Cellular responses of microcolonial rock fungi to long-term desiccation and subsequent rehydration. *Studies in Mycology* **61**: 91–97.
- Onofri S, Barreca D, Selbmann L, Isola D, Rabbow E, Horneck G, Vera JPP de, Hatton J and Zucconi L (2008). Resistance of Antarctic black fungi and cryptoendolithic communities to simulated space and Martian conditions. *Studies in Mycology* **61**: 99–109.
- Gueidan C, Ruibal Villaseñor C, Hoog GS de, Gorbushina AA, Untereiner WA and Lutzoni F (2008). A rock-inhabiting ancestor for mutualistic and pathogen-rich fungal lineages. *Studies in Mycology* **61**: 111–119.
- Deng S, Gerrits van den Ende AHG, Ram AFJ, Arentshorst M, Gräser Y, Hu H and Hoog GS de (2008). Evolution of *CDC42*, a putative virulence factor triggering meristematic growth in black yeasts. *Studies in Mycology* **61**: 121–129.
- Li DM, Hoog GS de, Lindhardt Saunte DM, Gerrits van den Ende AHG and Chen XR (2008). *Coniosporium epidermidis* sp. nov., a new species from human skin. *Studies in Mycology* **61**: 131–136.
- Vicente VA, Attili-Angelis D, Pie MR, Queiroz-Telles F, Cruz LM, Najafzadeh MJ, Hoog GS de, Zhao J and Pizzirani-Kleiner A (2008). Environmental isolation of black yeast-like fungi involved in human infection. *Studies in Mycology* **61**: 137–144.
- Sudhadham M, Prakitsin S, Sivichai S, Chaiyarat R, Dorrestein GM, Menken SBJ, and Hoog GS de (2008). The neurotropic black yeast *Exophiala dermatitidis* has a possible origin in the tropical rain forest. *Studies in Mycology* **61**: 145–156.

- Satow MM, Attili-Angelis D, Hoog GS de, Angelis DF and Vicente VA (2008). Selective factors involved in oil flotation isolation of black yeasts from the environment. *Studies in Mycology* **61**: 157–163.
- Rustler S, Chmura A, Sheldon RA and Stolz A (2008). Characterisation of the substrate specificity of the nitrile hydrolyzing system of the acidotolerant black yeast *Exophiala oligosperma* R1. *Studies in Mycology* **61**: 165–174.
- Badali H, Gueidan C, Najafzadeh MJ, Bonifaz A, Gerrits van den Ende AHG and Hoog GS de (2008). Biodiversity of the genus *Cladophialophora*. *Studies in Mycology* **61**: 175–191.